arXiv:0804.2239v2 [math-ph] 13 May 2008

Inverse Vector Operators

Shaon Sahoo [1]

Department of Physics, Indian Institute of Science, Bangalore 560012, India.

Abstract In different branches of physics, we frequently deal with vector del operator ($\vec{\nabla}$). This del operator is generally used to find curl or divergence of a vector function or gradient of a scalar function. In many important cases, we need to know the parent vector whose curl or divergence is known or require to find the parent scalar function whose gradient is known. But the task is not very easy, especially in case of finding vector potential whose curl is known. Here, 'inverse curl', 'inverse divergence' and 'inverse gradient' operators are defined to solve those problems easily. All operators are defined in the orthogonal curvilinear co-ordinate system.

Keywords del operator, curl, gradient, divergence, solenoidal vector, vector potentials, curvilinear co-ordinate system.

Here we consider u_1, u_2, u_3 as three mutually perpendicular families of co-ordinate surfaces and $\hat{e}_1$, $\hat{e}_2$, $\hat{e}_3$ as unit vectors normal to respective surfaces.

1. Inverse Curl Operator

At first we mention some symbols, used later.

a. $(\vec{\nabla}\times)^{-1}$ indicates inverse curl operator.

b. $(\int^+ + \int^-)(\partial u_1)$ does integration w.r.t. u_1, while other two variables of the integrand are to be treated fixed. If the integrand is the 3rd component (along $\hat{e}_3$) of a vector, then $\int^+(\partial u_1)$ acts only on the part of the integrand having the variable u_3. Similarly $\int^-(\partial u_1)$ acts only on the part of the integrand does not contain the variable u_3.

   For example, let $\phi(u_1,u_2,u_3) = \Psi_1(u_1,u_2) + \Psi_2(u_1,u_2,u_3)$ be the component of a vector along $\hat{e}_3$, then, $(\int^+ + \int^-)\phi\partial u_1 = \int^+ \Psi_2 \partial u_1 + \int^- \Psi_1 \partial u_1$ (u_2, u_3 are to be treated constant).

Now take two vectors $\vec{A} = A_1\hat{e}_1 + A_2\hat{e}_2 + A_3\hat{e}_3$ and $\vec{B} = B_1\hat{e}_1 + B_2\hat{e}_2 + B_3\hat{e}_3$, such that,

$$\vec{B} = \vec{\nabla}\times\vec{A} \tag{1}$$

Using inverse curl operator, this vector potential $\vec{A}$ can be expressed as,

$$\vec{A} = (\vec{\nabla}\times)^{-1}\vec{B} \tag{2}$$

---

[1] shaon@physics.iisc.ernet.in

From (1) it is obvious that,

$$\vec{\nabla}\cdot\vec{B}=\vec{\nabla}\cdot(\vec{\nabla}\times\vec{A})=0$$

$$\Rightarrow \frac{1}{h_1h_2h_3}[\frac{\partial}{\partial u_1}(h_2h_3B_1)+\frac{\partial}{\partial u_2}(h_3h_1B_2)+\frac{\partial}{\partial u_3}(h_1h_2B_3)]=0$$

$$\Rightarrow \frac{\partial}{\partial u_1}[c_1(u_1^+)+c_1(u_1^-)]+\frac{\partial}{\partial u_2}[c_2(u_2^+)+c_2(u_2^-)]+\frac{\partial}{\partial u_3}[c_3(u_3^+)+c_3(u_3^-)]=0$$

$$\Rightarrow \frac{\partial}{\partial u_1}c_1(u_1^+)+\frac{\partial}{\partial u_2}c_2(u_2^+)+\frac{\partial}{\partial u_3}c_3(u_3^+)=0 \qquad (3)$$

Where $c_1(u_1^+)$ and $c_1(u_1^-)$ are two parts of the term $h_2h_3B_1$, contains and does not contain u_1 respectively. Similarly, other terms can be defined.
Obviously $h_2h_3B_1=c_1(u_1^+)+c_1(u_1^-)$, and so on for other terms.
From (1) we get,

$$\vec{B}=\vec{\nabla}\times\vec{A}=\frac{1}{h_1h_2h_3}\begin{vmatrix} h_1\hat{e}_1 & h_2\hat{e}_2 & h_3\hat{e}_3 \\ \frac{\partial}{\partial u_1} & \frac{\partial}{\partial u_2} & \frac{\partial}{\partial u_3} \\ h_1A_1 & h_2A_2 & h_3A_3 \end{vmatrix}$$

$$\begin{aligned}\Rightarrow B_1\hat{e}_1+B_2\hat{e}_2+B_3\hat{e}_3=\ &\frac{\hat{e}_1}{h_2h_3}[\frac{\partial}{\partial u_2}(h_3A_3)-\frac{\partial}{\partial u_3}(h_2A_2)]\\ &+\frac{\hat{e}_2}{h_3h_1}[\frac{\partial}{\partial u_3}(h_1A_1)-\frac{\partial}{\partial u_1}(h_3A_3)]\\ &+\frac{\hat{e}_3}{h_1h_2}[\frac{\partial}{\partial u_1}(h_2A_2)-\frac{\partial}{\partial u_2}(h_1A_1)]\end{aligned}$$

Now, we have to choose h_1A_1, h_2A_2 and h_3A_3 in such a way that, r.h.s. of the above expression becomes similar to l.h.s.
Let (by inspection),

$h_1A_1=k_1\int c_2(u_2^+)\partial u_3+k_2\int c_3(u_3^+)\partial u_2+k_3\int c_2(u_2^-)\partial u_3+k_4\int c_3(u_3^-)\partial u_2$
Similarly,
$h_2A_2=k_5\int c_3(u_3^+)\partial u_1+k_6\int c_1(u_1^+)\partial u_3+k_7\int c_3(u_3^-)\partial u_1+k_8\int c_1(u_1^-)\partial u_3$

and, $h_3A_3=k_9\int c_1(u_1^+)\partial u_2+k_{10}\int c_2(u_2^+)\partial u_1+k_{11}\int c_1(u_1^-)\partial u_2+k_{12}\int c_2(u_2^-)\partial u_1$

where, $k_1,k_2,\cdots,k_{12}$ are some constants to be determined.
(Note that, c_1 terms are not included for h_1A_1, since, it does not appear in the expression of 1st component, B_1. So on for other two cases.)

Now, see the coefficient of $\hat{e}_1$ on the r.h.s. of the expression,

$$\frac{1}{h_1h_2}[\frac{\partial}{\partial u_1}(h_2A_2)-\frac{\partial}{\partial u_2}(h_1A_1)]$$

$$\begin{aligned}=\ &\frac{1}{h_1h_2}[\frac{\partial}{\partial u_1}\{k_5\int c_3(u_3^+)\partial u_1+k_6\int c_1(u_1^+)\partial u_3+k_7\int c_3(u_3^-)\partial u_1+k_8\int c_1(u_1^-)\partial u_3\}\\ &-\frac{\partial}{\partial u_2}\{k_1\int c_2(u_2^+)\partial u_3+k_2\int c_3(u_3^+)\partial u_2+k_3\int c_2(u_2^-)\partial u_3+k_4\int c_3(u_3^-)\partial u_2\}]\end{aligned}$$

$$\begin{aligned}=\ &\frac{1}{h_1h_2}[k_5c_3(u_3^+)+k_6\int\frac{\partial}{\partial u_1}c_1(u_1^+)\partial u_3+k_7c_3(u_3^-)+k_8\cdot 0-k_1\int\frac{\partial}{\partial u_2}c_2(u_2^+)\partial u_3\\ &-k_2c_3(u_3^+)-k_3\cdot 0-k_4c_3(u_3^-)]\end{aligned}$$

$$= \frac{1}{h_1h_2}[2k_5c_3(u_3^+) - k_1\int\{\frac{\partial}{\partial u_1}c_1(u_1^+) + \frac{\partial}{\partial u_2}c_2(u_2^+)\}\partial u_3 + 2k_7c_3(u_3^-)]$$

[Taking $k_5 = -k_2$, $k_7 = -k_4$ and $k_6 = -k_1$]

$$= \frac{1}{h_1h_2}[2k_5c_3(u_3^+) - k_1\int\{-\frac{\partial}{\partial u_3}c_3(u_3^+)\}\partial u_3 + 2k_7c_3(u_3^-)] \qquad \text{[Using eq}^n\text{(3)]}$$

$$= \frac{1}{h_1h_2}[2k_5c_3(u_3^+) + k_1c_3(u_3^+) + 2k_7c_3(u_3^-)]$$

$$= \frac{1}{h_1h_2}[3k_5c_3(u_3^+) + 2k_7c_3(u_3^-)] \qquad [\text{Taking } k_1 = k_5]$$

$$= \frac{1}{h_1h_2}[3\cdot\frac{1}{3}c_3(u_3^+) + 2\cdot\frac{1}{2}c_3(u_3^-)] \qquad [\text{Considering } k_5\text{=1/3 and } k_7\text{=1/2}]$$

$$= \frac{1}{h_1h_2}[c_3(u_3^+) + c_3(u_3^-)]$$

$$= \frac{1}{h_1h_2}(h_1h_2B_3)$$

$$= B_3$$

Similarly we can show that, if,

$k_1 = k_5 = k_9 = -k_2 = -k_6 = -k_{10} = 1/3$
and, $k_3 = k_7 = k_{11} = -k_4 = -k_8 = -k_{12} = 1/2$, then,

$\frac{1}{h_2h_3}[\frac{\partial}{\partial u_2}(h_3A_3) - \frac{\partial}{\partial u_3}(h_2A_2)] = B_1$ and $\frac{1}{h_1h_3}[\frac{\partial}{\partial u_3}(h_1A_1) - \frac{\partial}{\partial u_1}(h_3A_3)] = B_2$

So, using these values of $k_1, k_2, \cdots, k_{12}$, we get,

$$A_1 = \frac{1}{3h_1}[\int c_2(u_2^+)\partial u_3 - \int c_3(u_3^+)\partial u_2] + \frac{1}{2h_1}[\int c_2(u_2^-)\partial u_3 - \int c_3(u_3^-)\partial u_2]$$

$$A_2 = \frac{1}{3h_2}[\int c_3(u_3^+)\partial u_1 - \int c_1(u_1^+)\partial u_3] + \frac{1}{2h_2}[\int c_3(u_3^-)\partial u_1 - \int c_1(u_1^-)\partial u_3]$$

$$A_3 = \frac{1}{3h_3}[\int c_1(u_1^+)\partial u_2 - \int c_2(u_2^+)\partial u_1] + \frac{1}{2h_3}[\int c_1(u_1^-)\partial u_2 - \int c_2(u_2^-)\partial u_1]$$

Now from (2), we get,
$(\vec{\nabla}\times)^{-1}\vec{B} = \vec{A} = A_1\hat{e}_1 + A_2\hat{e}_2 + A_3\hat{e}_3$

$$\begin{aligned}
&= \frac{\hat{e}_1}{3h_1}[\int c_2(u_2^+)\partial u_3 - \int c_3(u_3^+)\partial u_2] + \frac{\hat{e}_1}{2h_1}[\int c_2(u_2^-)\partial u_3 - \int c_3(u_3^-)\partial u_2]\\
&\quad + \frac{\hat{e}_2}{3h_2}[\int c_3(u_3^+)\partial u_1 - \int c_1(u_1^+)\partial u_3] + \frac{\hat{e}_2}{2h_2}[\int c_3(u_3^-)\partial u_1 - \int c_1(u_1^-)\partial u_3]\\
&\quad + \frac{\hat{e}_3}{3h_3}[\int c_1(u_1^+)\partial u_2 - \int c_2(u_2^+)\partial u_1] + \frac{\hat{e}_3}{2h_3}[\int c_1(u_1^-)\partial u_2 - \int c_2(u_2^-)\partial u_1]
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\hat{e}_1}{3h_1}\begin{vmatrix}\int(\partial u_2) & \int(\partial u_3)\\ c_2(u_2^+) & c_3(u_3^+)\end{vmatrix} + \frac{\hat{e}_2}{3h_2}\begin{vmatrix}\int(\partial u_1) & \int(\partial u_3)\\ c_1(u_1^+) & c_3(u_3^+)\end{vmatrix} - \frac{\hat{e}_3}{3h_3}\begin{vmatrix}\int(\partial u_1) & \int(\partial u_2)\\ c_1(u_1^+) & c_2(u_2^+)\end{vmatrix}\\
&\quad - \frac{\hat{e}_1}{2h_1}\begin{vmatrix}\int(\partial u_2) & \int(\partial u_3)\\ c_2(u_2^-) & c_3(u_3^-)\end{vmatrix} + \frac{\hat{e}_2}{2h_2}\begin{vmatrix}\int(\partial u_1) & \int(\partial u_3)\\ c_1(u_1^-) & c_3(u_3^-)\end{vmatrix} - \frac{\hat{e}_3}{2h_3}\begin{vmatrix}\int(\partial u_1) & \int(\partial u_2)\\ c_1(u_1^-) & c_2(u_2^-)\end{vmatrix}
\end{aligned}$$

$$= -\frac{1}{3}\begin{vmatrix} \frac{\hat{e}_1}{h_1} & \frac{\hat{e}_2}{h_2} & \frac{\hat{e}_3}{h_3} \\ \int(\partial u_1) & \int(\partial u_2) & \int(\partial u_3) \\ c_1(u_1^+) & c_2(u_2^+) & c_3(u_3^+) \end{vmatrix} - \frac{1}{2}\begin{vmatrix} \frac{\hat{e}_1}{h_1} & \frac{\hat{e}_2}{h_2} & \frac{\hat{e}_3}{h_3} \\ \int(\partial u_1) & \int(\partial u_2) & \int(\partial u_3) \\ c_1(u_1^-) & c_2(u_2^-) & c_3(u_3^-) \end{vmatrix}$$

$$= -\begin{vmatrix} \frac{\hat{e}_1}{h_1} & \frac{\hat{e}_2}{h_2} & \frac{\hat{e}_3}{h_3} \\ \int(\partial u_1) & \int(\partial u_2) & \int(\partial u_3) \\ \frac{1}{3}c_1(u_1^+) + \frac{1}{2}c_1(u_1^-) & \frac{1}{3}c_2(u_2^+) + \frac{1}{2}c_2(u_2^-) & \frac{1}{3}c_3(u_3^+) + \frac{1}{2}c_3(u_3^-) \end{vmatrix}$$

$$= -\begin{vmatrix} \frac{\hat{e}_1}{h_1} & \frac{\hat{e}_2}{h_2} & \frac{\hat{e}_3}{h_3} \\ (\frac{1}{3}\int^+ + \frac{1}{2}\int^-)(\partial u_1) & (\frac{1}{3}\int^+ + \frac{1}{2}\int^-)(\partial u_2) & (\frac{1}{3}\int^+ + \frac{1}{2}\int^-)(\partial u_3) \\ h_2h_3B_1 & h_3h_1B_2 & h_1h_2B_3 \end{vmatrix}$$

So, this is the desired expression for our inverse curl operator.
Since, $\vec{\nabla} \times (\vec{\nabla}\phi) = 0$, so, it should be noted that, for a given solenoidal vector, there are many vector potentials, which can be in general expressed as, -

$$\vec{A} = (\vec{\nabla}\times)^{-1}\vec{B} + \vec{\nabla}\phi$$

where ϕ is an arbitrary scalar function.

One example in Cartesian system would clarify the procedure of getting vector potential from a solenoidal vector.
In this system, $\hat{e}_1$, $\hat{e}_2$, $\hat{e}_3$, u_1, u_2 and u_3 are respectively replaced by $\hat{i}$, $\hat{j}$, $\hat{k}$, x, y, and z. Here $h_1 = h_2 = h_3 = 1$.
Now take a solenoidal vector $\vec{B} = (xyz + y^2)\hat{i} + (xz + y)\hat{j} - z(1 + yz/2)\hat{k}$. (note $\vec{\nabla} \cdot \vec{B} = 0$)
So, vector potentials of $\hat{B}$ are,-

$$\vec{A} = (\vec{\nabla}\times)^{-1}\vec{B} + \vec{\nabla}\phi$$

$$= -\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ (\frac{1}{3}\int^+ + \frac{1}{2}\int^-)(\partial x) & (\frac{1}{3}\int^+ + \frac{1}{2}\int^-)(\partial y) & (\frac{1}{3}\int^+ + \frac{1}{2}\int^-)(\partial z) \\ xyz + y^2 & xz + y & -z - yz^2/2 \end{vmatrix} + \vec{\nabla}\phi$$

$$= -\hat{i}[\tfrac{1}{3}\int(-z - yz^2/2)\partial y + \tfrac{1}{2}\int 0\partial y - \tfrac{1}{3}\int y\partial z - \tfrac{1}{2}\int xz\partial z]$$

$$-\hat{j}[\tfrac{1}{3}\int xyz\partial z + \tfrac{1}{2}\int y^2\partial z - \tfrac{1}{3}\int(-z - yz^2/2)\partial x - \tfrac{1}{2}\int 0\partial x]$$

$$-\hat{k}[\tfrac{1}{3}\int y\partial x+\tfrac{1}{2}\int xz\partial x-\tfrac{1}{3}\int xyz\partial y-\tfrac{1}{2}\int y^2\partial y]+\vec{\nabla}\phi$$
$$=\hat{i}[(zy+z^2y^2/4)/3+(yz/3+xz^2/4)]-\hat{j}[(zx+yxz^2/2)/3+(xyz^2/6+y^2z/2)]$$
$$+\hat{k}[-(yx/3+x^2z/4)+xy^2z/6+y^3/6]+\vec{\nabla}\phi$$

Verification of the Result:

$$\vec{\nabla}\times\vec{A}=\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \begin{array}{c}(zy+z^2y^2/4)/3\\ +(yz/3+xz^2/4)\end{array} & \begin{array}{c}-(zx+xyz^2/2)/3\\ -(xyz^2/6+y^2z/2)\end{array} & \begin{array}{c}-(yx/3+x^2z/4)\\ +(xy^2z/6+y^3/6)\end{array} \end{vmatrix}+\vec{\nabla}\times(\vec{\nabla}\phi)$$

$$=\hat{i}(xyz+y^2)+\hat{j}(xz+y)-\hat{k}z(1+yz/2)+0$$
$$=\vec{B}$$

2. Inverse Divergence Operator

Here we indicate this operator by $(\vec{\nabla}\cdot)^{-1}$. Now if $\vec{A}$ and ϕ be a vector and a scalar function respectively, such that $\phi=\vec{\nabla}\cdot\vec{A}=\frac{1}{h_1h_2h_3}[\frac{\partial}{\partial u_1}(h_2h_3A_1)+\frac{\partial}{\partial u_2}(h_1h_3A_2)+\frac{\partial}{\partial u_3}(h_1h_2A_3)]$, then $\vec{A}$ can be expressed as $\vec{A}=(\vec{\nabla}\cdot)^{-1}\phi$,
We can here define the operator as (by inspection),
$(\vec{\nabla}\cdot)^{-1}=k_1\frac{\hat{e}_1}{h_2h_3}\int h_1h_2h_3(\partial u_1)+k_2\frac{\hat{e}_2}{h_3h_1}\int h_1h_2h_3(\partial u_2)+k_3\frac{\hat{e}_3}{h_1h_2}\int h_1h_2h_3(\partial u_3)$
where, $k_1+k_2+k_3=1$.

Justification:

For a given ϕ, $\vec{A}=(\vec{\nabla}\cdot)^{-1}\phi$

$$=k_1\frac{\hat{e}_1}{h_2h_3}\int h_1h_2h_3\phi(\partial u_1)+k_2\frac{\hat{e}_2}{h_3h_1}\int h_1h_2h_3\phi(\partial u_2)+k_3\frac{\hat{e}_3}{h_1h_2}\int h_1h_2h_3\phi(\partial u_3)$$

Now note that, divergence of the obtained vector gives the scalar function ϕ,

$$\vec{\nabla}\cdot\vec{A}=\frac{1}{h_1h_2h_3}[k_1h_1h_2h_3\phi+k_1h_1h_2h_3\phi+k_1h_1h_2h_3\phi]$$

$$=(k_1+k_2+k_3)\phi=\phi$$

Since, $\vec{\nabla}\cdot(\vec{\nabla}\times\vec{B})=0$, so, it should be noted that for a given scalar function, we can find many vectors, which can be expressed as,

$$\vec{A}=(\vec{\nabla}\cdot)^{-1}\phi+\vec{\nabla}\times\vec{B}$$

where $\vec{B}$ is any vector.

3. Inverse Gradient Operator

We indicate this operator by $(\vec{\nabla})^{-1}$

Let $\vec{A} = A_1\hat{e}_1 + A_2\hat{e}_2 + A_3\hat{e}_3$ and $\phi(u_1,u_2,u_3)$ be a vector and a scalar function, such that $\vec{A} = \vec{\nabla}\phi$.

So, vector field represented by $\vec{A}$ is conservative, thus, line integral of $\vec{A}$ is path independent. Choose (a,b,c) as initial point, such that $\phi(a,b,c)$ exists.

$$\text{Now } \phi(u_1,u_2,u_3) = \int_{(a,b,c)}^{(u_1,u_2,u_3)} d\phi \quad + c_0 \qquad [c_0 \text{ is integration constant}]$$

$$= \int_{(a,b,c)}^{(u_1,u_2,u_3)} (\vec{\nabla}\phi)\cdot\vec{dr} \quad + c_0$$

$$= \int_{(a,b,c)}^{(u_1,u_2,u_3)} \vec{A}\cdot\vec{dr} \quad + c_0$$

$$= \int_{(a,b,c)}^{(a,b,u_3)} A_3 h_3 du_3 + \int_{(a,b,u_3)}^{(a,u_2,u_3)} A_2 h_2 du_2 + \int_{(a,u_2,u_3)}^{(u_1,u_2,u_3)} A_1 h_1 du_1 \quad + c_0$$

So,
$\phi = (\vec{\nabla})^{-1}\vec{A}$

$$= \int_a^{u_1} A_1 h_1 \partial u_1 + \underset{u_1=a}{\int_b^{u_2} A_2 h_2 \partial u_2} + \underset{\substack{u_1=a\\ u_2=b}}{\int_c^{u_3} A_3 h_3 \partial u_3} + c_0$$

Here in the first integration u_1 is variable but other two (u_2,u_3) are fixed, and so on for other two integrations.

Note: The original work was done in 2002 and published in 2006 in Science and Culture [2]. This write-up is almost same as published one.

Reference

[1] Any text on vector del operator and curvilinear co-ordinate system. For example: by George B. Arfken and Hans J. Weber; *Mathematical Methods For Physicists*; Academic Press; 5th edition (2000); 1st and 2nd chapters.

[2] Shaon Sahoo, *Sci. & Cult.*, **72**, (1-2), 89 (2006).